

n-p Short-Range Correlations from (p,2p + n) Measurements

A. Tang^a, J. W. Watson^a, J. Aclander^b, J. Alster^b, G. Asryan^{d,e}, Y. Averichev^h, D. Barton^d, V. Baturin^{f,e},
N. Bukhtoyarova^{d,e}, A. Carroll^d, S. Heppelmann^f, A. Leksanov^f, Y. Makdisi^d, A. Malki^b, E. Minina^f, I. Navon^b,
H. Nicholson^g, A. Ogawa^f, Yu. Panebratsev^h, E. Piasezky^b, A. Schetkovsky^{f,e}, S. Shimanskiy^h, D. Zhalov^f

^a*Dept. of Physics, Kent State Univ., Kent, OH 44242, U.S.A.*

^b*School of Physics and Astronomy, Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 69978, Israel*

^c*Yerevan Physics Institute, Yerevan 375036, Armenia*

^d*Collider-Accelerator Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

^e*Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188350, Russia*

^f*Physics Department, Pennsylvania State University, University Park, PA 16801, U.S.A.*

^g*Dept. of Physics, Mount Holyoke College, South Hadley, MA 01075, U.S.A.*

^h*J.I.N.R., Dubna, Moscow 141980, Russia*

We studied the $^{12}\text{C}(\text{p},2\text{p}+\text{n})$ reaction at beam momenta of 5.9, 8.0 and 9.0 GeV/c. For quasielastic (p,2p) events we reconstructed $\mathbf{p}_f$ the momentum of the knocked-out proton before the reaction; $\mathbf{p}_f$ was then compared (event-by-event) with $\mathbf{p}_n$, the measured, coincident neutron momentum. For $|\mathbf{p}_n| > k_F = 0.220$ GeV/c (the Fermi momentum) a strong back-to-back directional correlation between $\mathbf{p}_f$ and $\mathbf{p}_n$ was observed, indicative of short-range n-p correlations. From $\mathbf{p}_n$ and $\mathbf{p}_f$ we constructed the distributions of c.m. and relative motion in the longitudinal direction for correlated pairs. After correcting for detection efficiency, flux attenuation and solid angle, we determined that 49 ± 13 % of events with $|\mathbf{p}_f| > k_F$ had directionally correlated neutrons with $|\mathbf{p}_n| > k_F$. Thus short-range 2N correlations are a major source of high-momentum nucleons in nuclei.

For the past half century, the dominant model for the structure of nuclei, especially light nuclei, has been the nuclear shell model. In the shell model, the long-range (~ 2 fm) part of the N-N force, in combination with the Pauli principle, produces an average potential in which the nucleons undergo nearly independent motion, and the residual interactions can be treated by perturbation theory. However the NN interaction is also highly repulsive at short-range (~ 0.4 fm). This short-range two-nucleon repulsion is responsible for the saturation density of nuclei, e.g. the nearly constant density in the interior of all stable nuclei heavier than ^4He . The effect of this two-nucleon short-range (2N-SRC) repulsion should also manifest itself in the motions of the nucleons in the nucleus. This topic has been actively pursued in recent years with electro-magnetic interactions. Recently, Aclander et al. [1] described a new technique for observing 2N-SRC with a high momentum transfer proton reaction, based on a proposal by Frankfurt and Strikman [2]. In this letter we show that this method not only demonstrates the presence of 2N-SRC but also provides quantitative information on the strength of 2N-SRC. We also obtained the first measured results of the c.m. momentum of the correlated NN pair as well as the relative momentum of the two nucleons in their c.m. frame.

For the quasi-elastic knockout of protons from nuclei, e.g. $^{12}\text{C}(\text{p},2\text{p})$ in [3] and for this work, we can use the Plane-Wave impulse approximation (PWIA) to reconstruct (event-by-event) the three momentum $\mathbf{p}_f$ that the struck target proton had before the reaction:

$$\mathbf{p}_f = \mathbf{p}_1 + \mathbf{p}_2 - \mathbf{p}_0 \quad (1)$$

where $\mathbf{p}_0$ is the momentum of the incident proton and $\mathbf{p}_1$ and $\mathbf{p}_2$ are the momenta of the two detected protons.

For p-p elastic scattering near 90° , the cross section $\frac{d\sigma}{dt}$ falls as s^{-10} where s and t are the standard Mandelstam variables for the square of the total c.m. energy and the squared momentum transfer, respectively. From this strong dependence on s (viz. s^{-10}) we expect p-p quasi-elastic scattering to occur preferentially with nuclear protons that have their momenta $\mathbf{p}_f$ roughly parallel to $\mathbf{p}_0$ the beam momentum, because this reduces s and increases $\frac{d\sigma}{dt}$ [4]. When two nucleons in a nucleus interact at short range, they must have large, nearly equal, momenta in opposite directions (because of their strong repulsion at short range) [5]. If the pair is broken by a quasi-elastic interaction one of the nucleons may emerge from the nucleus, with relatively large momentum, in a direction opposite to original momentum of its partner. Our experiment then consists of measuring the two protons of the quasi-elastic scattering in triple coincidence with the emerging correlated neutron. Since $\mathbf{p}_f$ is roughly parallel to the beam, we placed 36 neutron detectors primarily in the backward hemisphere, to search for neutrons with $\mathbf{p}_n \approx -\mathbf{p}_f$. The experimental details were described in [6].

The measurement reported here (and in [1]) were taken with the EVA spectrometer [6,7] at the AGS accelerator at Brookhaven National Laboratory. EVA consists of a 0.8 T superconducting solenoid, 3.3 m long and 2 m in diameter, with the beam incident along the central axis. Coincident pairs of high transverse-momentum ($\mathbf{p}_t$) protons are detected with four concentric cylinders of straw-tube drift chambers. EVA was designed for studying

quasi-elastic (p,2p) reactions near 90° c.m. and to investigate nuclear transparency [8,9].

For all of the results discussed below we applied the four following cuts to the data:

- (1). The projectile should be a proton, as determined by Čerenkov detectors in the beam.
- (2). There should be two (and only two) high- p_t (> 0.6 GeV/c) positive-charge tracks.
- (3). The missing energy, E_{miss} should be appropriate for quasi-elastic scattering, and was defined by the resolution at each energy [3].
- (4). The neutron momentum should be in the range: $0.05 < p_n < 0.55$ GeV/c.
- (5). The laboratory azimuthal angles for the two detected protons must lie within $\pm 45^\circ$ of the plane through EVA parallel to the face of the active neutron detector.

Cut (5) was implemented so that we could combine our results with those in [1], which were also taken with EVA but with more limited acceptance.




FIG. 1. Scatter plot of p_f^{up} vs. p_n with cuts 1, 2, 3, 4 and 5 (see text) for $^{12}\text{C}(p,2p+n)$ events. Data labelled “98” (solid symbols) are for this work. Data labelled “94” are from Aclander, et al. (Ref. [1]). The vertical dashed line at 0.22 GeV/c corresponds to k_F , the Fermi momentum for ^{12}C .

Our primary objective was to identify short-range correlated n-p pairs in ^{12}C nuclei and to determine their properties. The signature of such pairs is that the momentum of the detected neutron, $\mathbf{p}_n \approx -\mathbf{p}_f$. We should expect, however, that the motion of the c.m. of such an n-p pair in the nucleus will keep this from being a perfect equality. The simplest directional correlation we can construct is “up-down”. The neutron detectors were all

placed below the mid-plane of EVA, so the neutrons were all detected in the “downward” direction. We can then compare this with the “upward” component of $\mathbf{p}_f$.

Figure 1 is a scatter plot for (p,2p+n) events of p_f^{up} vs. the neutron momentum p_n , where p_f^{up} is the projection of $\mathbf{p}_f$ on an axis normal to the faces of the neutron detectors. The vertical dashed line in Fig. 1 is at $k_F = 0.22$ GeV/c, the Fermi momentum for ^{12}C [10]. Similar to [3], the experimental uncertainties in Fig. 1 are derived from resolutions measured for p-p elastic scattering. We note in Fig. 1, that there is a striking difference in the distribution of p_f^{up} for $p_n < k_F$ and $p_n \geq k_F$. For $p_n < k_F$ the events are distributed nearly equally between those with $p_f^{up} > 0$ (“up”) and $p_f^{up} < 0$ (“down”). By contrast, for $p_n \geq k_F$, a large fraction of the events have $p_f^{up} > 0$. Thus $\mathbf{p}_f$ is predominantly “up” when $\mathbf{p}_n$ is (by definition) “down” for $p_n \geq k_F$. This is a strong confirmation of what was reported by Aclander et al. in [1].

We now construct the full directional correlation between $\mathbf{p}_f$ and $\mathbf{p}_n$ as

$$\cos\gamma = \frac{\mathbf{p}_f \cdot \mathbf{p}_n}{|\mathbf{p}_f| |\mathbf{p}_n|} \quad (2)$$

where γ is the angle between $\mathbf{p}_f$ and $\mathbf{p}_n$.




FIG. 2. Plots of $\cos\gamma$, where γ is the angle between $\mathbf{p}_n$ and $\mathbf{p}_f$, for $^{12}\text{C}(p,2p+n)$ events. Panel (a) is for events with $p_n > 0.22$ GeV/c, and panel (b) is for events with $p_n < 0.22$ GeV/c; 0.22 GeV/c = k_F , the Fermi momentum for ^{12}C .

In Fig. 2a and 2b we plot $\cos\gamma$ for $p_n \geq k_F = 0.22$ GeV/c and $p_n < k_F$, respectively. As in Fig. 1 there is a pronounced difference between the distributions for $p_n \geq k_F$ and $p_n < k_F$. Fig. 2a shows a strong “back-to-back” directional correlation with the distribution peaking at $\cos\gamma = -1$; only 3 events have $\cos\gamma > 0$. By contrast, in Fig. 2b, the distribution is nearly uniform in $\cos\gamma$, and

the numbers of events for $\cos\gamma < 0$ and $\cos\gamma > 0$, are the same within statistics (50 and 40, respectively). For Fig. 2a ($p_n > k_F$), the probability that 41 uncorrelated events could be distributed 38 to 3 is vanishing small ($\sim 10^{-8}$). To check the results in Fig. 2, we generated background spectra of events with two high p_t tracks plus one soft track in EVA. The resulting spectra for both $p_n > k_F$ and $p_n < k_F$ were small and flat, indicating that the peak at $\cos\gamma = -1$ in Fig. 2a is not an instrumental artifact, and that backgrounds are small.

The effects of initial-state and final-state interactions (ISIs and FSIs) are expected to be larger in the transverse than in the longitudinal direction. Therefore, in extracting the relative and c.m. motion for correlated n-p pairs we focus on the longitudinal (z) components of $\mathbf{p}_n$ and $\mathbf{p}_f$. For beams of high-energy protons, the natural variables for representing the motion of a nucleon in the nucleus are the light-cone variables $\mathbf{p}_t$ and $\alpha = (E - p_z)/m$. Note that $\alpha \approx 1$ for $p_z = 0$, therefore, we would expect $\alpha_f + \alpha_n \approx 2$ for a correlated pair with $\mathbf{p}_n \approx -\mathbf{p}_f$. ($\alpha_f = (E - p_{fz})/m$; $\alpha_n = (E - p_{nz})/m$)




FIG. 3. Plots of the light-cone variables α_p , α_n , and their sum $\alpha_p + \alpha_n$ for events with $p_n > k_F = 0.22$ GeV/c. Data are for $^{12}\text{C}(p, 2p+n)$ events. Each event has been “ s -weighted”, as described in the text.

As noted earlier, the p-p cross section, $\frac{d\sigma}{dt}$, near 90° c.m. is proportional to s^{-10} . We therefore need to correct for this “reaction bias” when looking at longitudinal variables. The differential cross section as a function of the solid angle, $\frac{d\sigma}{d\Omega}$, is related to $\frac{d\sigma}{dt}$ as:

$$\frac{d\sigma}{dt} = \frac{4\pi}{(s - 4m^2)} \frac{d\sigma}{d\Omega} \quad (3)$$

Therefore, for large s , $\frac{d\sigma}{d\Omega}$ is approximately proportional to s^{-9} , and this strong s -dependence enhances quasielastic reactions with low s for the p-p collisions. Protons in the nucleus with longitudinal momentum in the same direction as the beam are thus more likely to be knocked

out. We therefore weighted each event by a correction factor equal to $(s/s_0)^9$, to obtain the nuclear distributions without the reaction bias, where s_0 is the total c.m. energy for pp $\rightarrow$ pp at each beam momentum, and s is calculated for each event from $\mathbf{p}_f$ for that event.

In Figure 3 we show plots of α_f , α_n and $\alpha_f + \alpha_n$ (all with s -weighting) for events with $p_n > k_F$. We note for Fig. 3 that α_f (for the proton) is generally < 1 and $\alpha_n > 1$. Of course, our placement of the neutron detectors primarily in the backward hemisphere forces α_n to be largely > 1 . In Fig. 3, the spread of $\alpha_n + \alpha_f$ about 2 should be due to the c.m. motion of the pair. Cioffi degli Atti *et al.* [11] emphasized the importance of the c.m. motion of correlated pairs for explaining nucleon spectral functions at large momenta and removal energies.

In the longitudinal direction:

$$p_z^{cm} = p_{nz} + p_{fz}. \quad (4)$$

By approximating $E_p \approx E_n \approx m$, we obtain

$$\alpha_p + \alpha_n = \left(1 - \frac{p_{fz}}{m}\right) + \left(1 - \frac{p_{nz}}{m}\right) \quad (5)$$

which leads to

$$p_z^{cm} = 2m\left(1 - \frac{\alpha_p + \alpha_n}{2}\right). \quad (6)$$

The longitudinal momentum of the particles in their c.m. frame can be extracted from the difference of the α variables. By again approximating $E_p \approx E_n \approx m$, we obtain

$$\alpha_p - \alpha_n = \left(1 - \frac{p_{fz}}{m}\right) - \left(1 - \frac{p_{nz}}{m}\right) = \frac{p_{nz} - p_{fz}}{m}, \quad (7)$$

which leads to

$$p_z^{rel} = m|\alpha_p - \alpha_n|. \quad (8)$$




FIG. 4. Plots of (a) p_z^{cm} and (b) p_z^{rel} for correlated n-p pairs in ^{12}C , for $^{12}\text{C}(p, 2p+n)$ events. Each event has been “ s -weighted”, as described in the text.

Figure 4a and 4b are plots of p_z^{cm} and p_z^{rel} . For the data for p_z^{cm} in Fig. 4a, the centroid is -0.013 ± 0.027 GeV/c. The spread in the distribution is $\sigma = 0.143 \pm 0.017$ GeV/c. For the data for p_z^{rel} in Fig. 4b, the centroid is 0.289 ± 0.017 GeV/c, and the width is $\sigma = 0.097 \pm 0.007$ GeV/c.

An interesting number which can be extracted from our data, is the fraction of $^{12}\text{C}(p,2p)$ events which have correlated neutrons with $\mathbf{p}_n \approx -\mathbf{p}_f$ when $p_n, p_f \geq k_F$. To extract this number, we need to correct the measured neutron flux for neutron detection efficiency and flux attenuation, and for solid-angle coverage. Our neutron detectors were placed almost entirely in the backward hemisphere, so we calculate the fraction of the 2π solid angle for the backward hemisphere covered by our detectors.

What we then calculated was:

$$F = \frac{\text{corrected \# of } (p,2p+n) \text{ events}}{\text{\# of } (p,2p) \text{ events}} = \frac{A}{B} \quad (9)$$

for the same data sample. B was obtained by applying cuts (1), (2), (3) and (5), to events with $p_f \geq k_F$ for our 5.9 GeV runs excluding the data reported in [1]. The quantity B = 2205 then was all events satisfying the above cuts and data selection. The quantity A was obtained from the sample of all 18 (p,2p+n) events in the sample B with $p_n \geq k_F$, where a correction for flux attenuation (t) [12] and detection efficiency (ϵ) [13] was applied event-by-event. The resulting quantity was then corrected for the solid-angle coverage to obtain A:

$$A = \frac{2\pi}{\Delta\Omega} \sum_{i=1}^{18} \frac{1}{\epsilon_i} \cdot \frac{1}{t_i} = 1090. \quad (10)$$

The average value of $\epsilon_i^{-1} t_i^{-1}$ was 8.2 ± 0.82 and $2\pi/\Delta\Omega = 7.42$. We then obtain

$$F = \frac{A}{B} = 0.49 \pm 0.13. \quad (11)$$

The sample of 18 measured (p,2p+n) events is clearly small, and the uncertainty in F is determined largely by this sample size. The result is still compelling: roughly half of the measured quasielastic (p,2p) events with $p_f > k_F$ have a neutron emitted in the backward hemisphere with $p_n > k_F$. We note that this result is very similar to that reported by Malki, et al. [6] for hard inclusive (p,2p+n) measurements.

In summary, for quasi-elastic (p,2p) events we reconstructed $\mathbf{p}_f$, the momentum that the struck proton had in ^{12}C before the reaction. Then, for neutrons with momenta $p_n > 0.220$ GeV/c (which is the k_F , Fermi momentum for ^{12}C), we found a strong directional correlation between $\mathbf{p}_f$ and $\mathbf{p}_n$, namely $\mathbf{p}_n \approx -\mathbf{p}_f$. By contrast, for $p_n < k_F$ we found no correlation in directions. This was evident in the one-dimensional “up-down” and longitudinal correlations and the full three-dimensional directional correlation. For the longitudinal direction, where

ISI and FSI effects should be small, we extracted the distributions of c.m. and relative motion for n-p pairs.

We conclude, therefore, that neutrons emitted into the backward hemisphere with $p_n > k_F$ come from n-p SRC, since SRC is a natural mechanism to explain such momentum-correlated pairs. An analysis of the sample of events for 5.9 GeV/c beam momentum with $p_f > k_F$, indicates that $49 \pm 13\%$ of these events have a correlated neutron with $p_n > k_F$. Because this measured fraction includes only n-p and not p-p SRC, the total correlated fraction must be even larger. Therefore we conclude that 2N SRC must be a major source of high-momentum nucleons in nuclei. We also measured the longitudinal components of the c.m. momentum of the correlated pn pair and the relative momentum of the pn pair in its c.m. system. Further applications of the technique used in this work and in Ref. [1] are planned [14] with the high-intensity c.w. electron beams at the Thomas Jefferson National Accelerator Facility.

ACKNOWLEDGMENTS

We are pleased to acknowledge the assistance of the AGS staff in building and rebuilding the detector and supporting the experiment, particularly our liaison engineer C. Pearson, and the continuing support of Drs. D. Lowenstein and P. Pile. This research was supported by the U.S. - Israel Binational Science foundation, the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities, NSF grants No. PHY-9501114, PHY-9722519, PHY-0099387 and the U.S. Department of Energy grant No. DEFG0290ER40553.

-
- [1] J. Aclander *et al.*, Phys. Lett. B **453**, 211 (1999).
 - [2] L.L. Frankfurt and M.I. Strikman, Phys. Rep. **76**, 214 (1981); Phys. Rep. **160**, 235 (1988).
 - [3] Y. Mardor *et al.*, Phys. Lett. B **437**, 257 (1998).
 - [4] G.R. Farrar *et al.*, Phys. Rev. Lett. **62**, 1095 (1989).
 - [5] A. De Shalit and H. Feshbach, Theoretical Nuclear Physics, Wiley, 1974.
 - [6] A. Malki *et al.*, Phys. Rev. C **65**, 015207 (2001).
 - [7] J.Y. Wu *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **349**, 183 (1994).
 - [8] I. Mardor *et al.*, Phys. Rev. Lett. **81**, 5085 (1998).
 - [9] A. Leksanov *et al.*, Phys. Rev. Lett. **87**, 212301 (2001).
 - [10] E.J. Moniz *et al.*, Phys. Rev. Lett. **26**, 445 (1971).
 - [11] C. Cioffi degli Atti, S. Simula, L.L. Frankfurt and M. Strikman, Phys. Rev. C **44**, 7 (1991).
 - [12] D.J. Hughes and R.B. Schwartz, BNL report 325 (1958).
 - [13] R.A. Cecil *et al.*, Nucl. Inst. and Meth. **161**, 439 (1979).
 - [14] TJNAF experiment E01-015, E. Piasetzky, W. Bertozzi, J. W. Watson, and S. Wood spokespersons (2001).